



NR 32% BEEF CATTLE SUPPLEMENT PELLETTED MEDICATED

Medicated with monensin sodium 356 mg/kg as an aid in the prevention of Coccidiosis caused by *E. bovis* and *E. zuernii* in beef cattle.
A list of the ingredients used in the manufacture of this feed may be obtained from the manufacturer or registrant.
This feed contains added selenium at 4.5 mg/kg.

GUARANTEED ANALYSIS:

Crude Protein, Min	32.00 %
ECP from NPN*, Max	20.00 %
Crude Fibre, Max	15.00 %
Crude Fat, Min	1.50 %
Calcium, Act	7.00 %
Phosphorus, Act	0.56 %
Magnesium, Act	1.00 %
Salt, Act	5.09 %
Sodium, Act	2.00 %
Iodine, Act	15 mg/kg
Cobalt, Act	6 mg/kg
Iron, Act	1,003 mg/kg
Copper, Act	400 mg/kg
Manganese, Act	622 mg/kg
Zinc, Act	1,500 mg/kg
Vitamin A, min	150,000 iu/kg
Vitamin D ³ , min	15,000 iu/kg
Vitamin E, min	450 iu/kg
Flourine, Max	50 mg/kg

*Equivalent Crude Protein from Non-Protein Sources

MIXING DIRECTIONS: Mix 100 kilograms of 32% Beef Cattle Supplement Pelleted with 900 kilograms of barley and 1 kg Dyna-K.

FEEDING DIRECTIONS:

Feed the ration made from this supplement to beef cattle at a rate of 3.65 kg to beef cattle weighing 228 kg and consuming 2.5 % of body weight on a dry matter basis. Always have a source of clean water with adequate amounts available at all times. Provide free choice good quality forage. Have free choice salt available at all times. Feeding directions must be followed carefully. Do not feed undiluted. Contact a New Rosedale Feedmill representative for more information.

WEIGHT OF BEEF CATTLE KG	DRY MATTER INTAKE KG	FEEDING RATE KG
228	5.70	3.650
300	7.50	4.815
350	8.75	5.618
400	10.00	6.420

WARNING:

1. Do not supplement monensin from other sources (e.g., other feedstuffs containing monensin or slow release devices containing monensin).
2. No withdrawal period or milk withholding time is required when treated at the recommended dose of 22 mg monensin per kg (0.0022%) of complete diet.
3. When mixing and handling monensin, avoid inhalation, oral exposure and direct contact with skin or eyes. Use protective clothing, impervious gloves and dust mask. Operators should wash thoroughly with soap and water after handling.
4. Keep out of reach of children.

CAUTION:

1. Do not exceed recommended levels as reduced average daily gains may result.
2. Do not allow dogs, horses, or other equines or guinea fowl access to formulations containing monensin. Ingestion of monensin by these species has been fatal.
3. Do not use monensin-medicated feed for the treatment of outbreaks of coccidiosis.
4. May be used in feeds containing the following pellet-binding agents: bentonite (2%), attapulgit (2%), kaolin (2.5%), lignin sulfonate (4%), carboxymethylcellulose (0.1%), or Agri-Colloid.
5. Directions for use must be carefully followed.
6. Do not use in association with another feed containing supplemental selenium.

Individual results from the use of this product may vary due to management, environmental, genetic, health and sanitation differences; therefore, New Rosedale Feed Mill does not guarantee individual results.

Manufactured by:
NEW ROSEDALE FEED MILL
PORTAGE LA PRAIRIE, MANITOBA

Net weight on bulk invoice or 25 kg per bag when bagged.

Formula Code: 4108P